

PRODUCT INFORMATION

Product Name : Competent Cell DH5 α
Code No. : DS220
Size : 100 μ l \times 10
Competency : $> 5 \times 10^8$ cfu/ μ g (pUC19)
Supplied product : SOC medium, 1 ml \times 10

This product is for research use only

Description :

Competent Cells DH5 α is a high-efficiency chemically competent cell from *E. coli* DH5 α strain (one of the standard strains for molecular biology research) and suitable for a wide variety of cloning applications. The DH5 α cell has mutation of ϕ 80*lacZ* Δ M15 and lacks *laqI*^q gene, which allows blue-white color screening of transformants with X-gal (IPTG is not required).

Genotype of *E. coli* strain DH5 α :

*supE44, Δ lacU169(ϕ 80*lacZ* Δ M15), *hsdR17, recA1, endA1, gyrA96, thi-1, relA1**

Quality Control :

Transformation was carried out according to the method described in this Product Information using supercoiled pUC19 plasmid. Transformants were plated on LB plates containing 50 μ g/ml ampicillin. The efficiency was confirmed to be greater than 5×10^8 cfu/ μ g.

Storage condition :

Stable at -80°C with little or no loss in transformation efficiency for 12 months from the date of receipt. Competent Cells are sensitive to variation in temperature. Must be stored at -80°C. Upon receipt, store the Competent Cell DH5 α in a freezer at -80°C directly from a dry ice shipping box and store SOC medium at room temperature or at -80°C.

Handling of competent cells :

- Competent cells are sensitive to mechanical shock. Excessive mixing should be avoided.
- After thawing competent cells on ice, cells tend to lose transformation efficiency gradually. Transformation should be started immediately following thawing cells on ice.
- Use of refrozen competent cells is not recommended.

Composition of SOC medium supplied :

20 g/L	tryptone
5 g/L	yeast extract
0.5 g/L	NaCl
0.186 g/L	KCl
2.4 g/L	MgSO ₄ , anhydrous
4 g/L	glucose

PRODUCT INFORMATION

Transformation Procedure :

- Materials to be supplied by user
 - LB plates with antibiotic
 - Ice bucket with ice
 - 15 ml sterilized-polypropylene culture tubes
 - 42°C water bath
 - 37°C shaker
 - Sterile spreaders
 - 37°C incubator

If blue-white screening is required to select transformants,

- 20 mg/ml X-Gal in dimethylformamide (DMF)

● Transformation

1. Thaw one tube of competent cells on ice. One tube contains 100 µl of cells for each transformation.
2. Add DNA sample* directly into the competent cells and mix by flicking the tube.

* The volume of DNA sample should not exceed 5 % of that of competent cells (i.e. for 100 µl of competent cells, use ≤ 5 µl).

3. Incubate the tube on ice for 20 minutes.
4. Heat Shock the cells by placing the tube in 42°C water bath for 45 seconds. Do not mix or shake.
5. Remove the tube from the 42°C bath and place it on ice for 2 min.
6. Transfer the cells to a 15 ml sterilized culture tube containing 0.9 ml of SOC medium (pre-warmed at room temperature to 37°C). Culture the cells at 37°C for 1 hr in a shaker.
7. Spread an aliquot of the cells onto an LB agar plate containing appropriate antibiotic.
If blue-white color screening is required, spread 25 µl of 20 mg/ml X-Gal onto an LB agar plate and allow the reagent to absorb 30 minutes prior to inoculating cells. As DH5α does not have *lacI^q*, IPTG is not required for blue-white screening.
8. Incubate the plate at 37°C overnight.

Reference:

Sambrook, J. and Russell, D.W. (2001) Molecular Cloning: A Laboratory Manual, 3rd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

Related Products:

DS225	Jet Competent Cell (DH5α)	DS210	Competent Cell JM109
DS218	Electrocompetent Cell JM109	DS228	Electrocompetent Cell DH5α
DS229	Giga Competent Cell (DH5α)		